

Department of Agriculture, Fisheries and Forestry ABARES

Agricultural monitoring in Australia using satellite data: status and developments



Kate Harle, Lucy Randall, Alex Held (CSIRO), Medhavy Thankappan (Geoscience Australia)

21 November 2013




Australian agriculture: snapshot

- 59% land area and 52% national water use (2009–10)
- 2% GDP (2011-12) A\$48 billion
- Export around 75% (2011-12)
 - crops & livestock = A\$36 billion

 2.1% global exports (agriculture, fisheries & forestry -2012)

10% of global wheat exports (2005-2010)

- Challenges:
 - Land degradation
 - Variable climates (and climate change)
 - Increasing productivity to meet global food demand



2007 US\$10 billion

Source: ABARES 2012


Satellite data for agricultural monitoring: key uses



- Monitoring land condition
- Within season forecasting
- Extreme events early warning and damage assessment
- Validation and calibration efforts
- ➤ State and Federal governments, industry, research agencies e.g. ABARES, CSIRO, Geoscience Australia, Terrestrial Ecosystem Resource Network (TERN), GEOGLAM

National Ground Cover Monitoring project

 Nationally validated remotely sensed fractional cover - MODIS (non-woody and litter near soil surface)



National Ground Cover Monitoring project

- Nationally validated remotely sensed fractional cover (non-woody and litter near soil surface)
- Ground cover monitoring over large spatial extents at multiple scales through time
- Assessment of environmental targets related to soil erosion and land management
- Using satellite data to track:
 - change in ground cover compared to previous years influence of management practices
 - where cover persistently low potential for improvement?
 - wind erosion risk (threshold 50% bare soil)
 - water erosion risk (threshold of 70% bare soil)



Photo: John Leys, NSW OEH

Landsat time series

- Archive of calibrated time series of images (1998 to 2012) –
 Geoscience Australia
- Data set land cover change, water inundation etc.
- Available on National Computing Infrastructure, CRC for Spatial Information, TERN
- Developing web portal for access and research






Products for farmers

Web decision making tools using satellite data

- e.g. Pastures from Space®
 - Whole-farm & within-paddock tactical & strategic decision support
 - Pasture growth rate (kg/ha/day)
 - Pasture biomass (kg/ha) calibrated at intensive sites
 - Spatial and historical regional comparisons
 - Pasture growth model uses Modis NDVI + radiation, climate & soils data






Within season forecasting

- Focus on crop forecasting (wheat and sorghum)
- Satellite data used to improve crop models
 - Green up (satellite data)
 - Other data used climate and soil moisture
- Satellite data used to monitor crop condition through season
 - NDVI anomalies
 - Bureau of Meteorology
 - GEOGLAM
 - GEOGLAM Rangelands initiative NDVI, fractional cover, models
- Integrated crop forecasting (experimental) complement crop forecasting using survey based techniques









- Monitor global lands integral to producing animal protein 'free-range', open-field basis.
- Global & regional population information beef cattle, goats, sheep, camels, pork, dairy cattle, buffalo & deer.
- Excludes fisheries, intensive livestock cattle yards, piggeries, poultry
- NDVI, fractional cover and models
- Activities
 - International workshop Sydney October 2013
 - Implementation plan finalised Jan 2014
 - France 2nd meeting April 2014
 - Kenya 2nd half 2014 International Livestock Research Institute

Early warning and damage assessment

- Fire Sentinel, Landgate Firewatch, Northern Australia Fire Information
- Drought
- Flood
- Plague locusts



Landgate Firewatch


Sentinel – fire hot spots



Flooding central QLD Jan 2011




Note: Dark blue areas indicate water. Light blue and white are clouds.



Future development – Australian context

- Integrated Crop Forecasting
 - Developing fully functional system
 - Incorporating longer term climate forecasts
- Improve identification of crop area
 - crop identification
 - improve determination of green up
- Coping with cloudy data coastal & monsoonal areas
- Better coordination and access to existing & new data
 - New satellites
 - Soil moisture
 - Climate data (historical and seasonal forecast)
- GEOGLAM rangelands



Thank you




Kate Harle, Lucy Randall, Alex Held (CSIRO), Medhavy Thankappan (Geoscience Australia)

21 November 2013

Assessment of wind erosion risk

Counts the number of time periods in a season that bare soil:


- Exceeds 50% (wind erosion)



Products for farmers

Web decision making tools using satellite data

- > e.g. VegMachine
 - Regional monitoring and data
 - Resource condition, carrying capacity, grazing impact
 - Spatial and historical regional comparisons within local area
 - Landsat cover index trends, climate
 & soils data



Wallace et al 1994, 1998, 2006 LCCA – operational scale Karfs et al 2000, 2002...

Slide 14

KH1 check this

check this Katherine Harle, 11/15/2013