

IAC-11-B5.2.9

USING THE DSST SEMI-ANALYTICAL ORBIT PROPAGATOR PACKAGE VIA THE
NONDYWEBTOOLS/ASTRODYWEBTOOLS OPEN SCIENCE ENVIRONMENT

Juan F. San-Juan

Universidad de La Rioja, Logroño, Spain, juanfelix.sanjuan@unirioja.es

Martín Lara

Real Observatorio de la Armada, San Fernando, Spain, mlara@roa.es

Rosario López

Universidad de La Rioja, Logroño, Spain, rolopeg@gmail.com

Luis M. López

Universidad de La Rioja, Logroño, Spain, luis-maria.lopezo@unirioja.es

Zachary J. Folcik

Massachusetts Institute of Technology, Cambridge, USA, folcikz@gmail.com

Brian Weeden

Secure World Foundation, Washington, DC, USA, bweeden@swfound.org

Paul J. Cefola

University at Buffalo SUNY, Amherst, New York, USA, paulcefo@buffalo.edu

An initiative within the realm of Space Surveillance Awareness (SSA) is to create an open source software suite that can provide all space actors access to the basic SSA analysis tools needed to operate safely and efficiently in space. These applications include observation compression, orbit propagators, state transition matrix, weighted least squares, improved nonlinear Kalman filters, realistic process noise, atmosphere density variations, observation data association and observation data simulation which will be rewritten in a modern distributed object-oriented computing environment. It is the intent to make these applications standable via an Internet framework. While the project is being developed, one of the tools considered, the DSST standalone, which is an accurate Semi-Analytical Satellite Theory, can be used through Internet framework Astrody^{Web}_{Tools}. This provides an Astrodynamics framework so as to carry out open science in which specialized applications can be integrated, and to encourage scientific collaboration through Internet. This framework, through a user-friendly web interface, allows the user to choose applications, introduce data and select appropriate constraints in an intuitive and easy way, with the help of graphical interface options. After that, the application is executed in real-time and the critical information about program behavior and output (graphical representation of data, statistical analysis or whichever manipulation therein) is shown via the same web interface and can be downloaded to users' computers.

I. INTRODUCTION

Hundreds of thousands of human-made objects are orbiting at different altitudes around Earth. These objects include active and inactive satellites, spent rocket bodies used to launch satellites, and a wide variety of objects of all shapes and sizes, which have been released during missions or have come from collisions, explosions, and so on. Tracking them as precisely as possible, and analyzing their potential impact on space activities, is an essential part of Space Surveillance Awareness (SSA).

The accuracy of the SSA orbital data products are affected by the evolution of the sensors feeding the network, by the knowledge and control of the errors in

the sensor network, by the knowledge of the space environment, by the available computing resources (both hardware and software), and by the number of space objects to be monitored. However, the quality and quantity of the orbital data products available to space operators has evolved slowly over time. Further, error analysis of key issues is still in flux. The Iridium /Cosmos collision event in 2009 demonstrated that accurate SSA data was not sufficiently integrated into owner/operator management of satellite missions¹. Further, lack of data fusion and inadequate modeling of the satellite motion are technical factors contributing to erroneous orbit prediction of the close approach distances in many instances.

In 2010, P. J. Cefola, B. Weeden and C. Levit² presented a plan for an “Open Source Software Suite for Space Situational Awareness and Space Object Catalog Work.” This paper addressed two major issues:

1. the problem of adapting legacy SSA software tools to modern computing environments
2. the addition of the new analytic functionality to the operational SSA toolbox

Adapting complex scientific software to modern computing environments includes:

- non-invasive encapsulation of legacy binaries
- migration of SSA tools to a language platform employing object-oriented and component technologies such as C++
- a plan for creating a Web 2.0 architecture

Observation compression, certain orbit propagators, improved nonlinear Kalman Filters all are examples of 'new functionality' for the operational SSA environment.

For example, this project currently intends to include a variety of orbit propagators: Numerical Integration (Special Perturbations), DSST³, Brouwer-Lyddane⁴, NORAD GP⁵ (SGP, SGP4, SGP8, with tesseral m-daily option), NORAD HANDE⁶ (with tesseral m-daily option) NAVSPASUR PPT^{7,8} (with tesseral m-daily option), and the Russian A, AP, and NA theories⁹.

While the re-engineering processes of these orbit propagators is being carried out, at the same time we are working on their integration into a Web framework, which will allow the scientific community free use of these tools through the Internet.

One of these orbit propagators is DSST (Draper Semi-Analytical Satellite Theory) which was developed by P. J. Cefola, W. McClain, L. Early, R. Proulx, M. Slutsky and their colleagues at the Computer Sciences Corporation and the Charles Draper Laboratory (CSDL) in the 1970s and 1980s. In its development at the CSDL, DSST also benefited from numerous enhancements made by Massachusetts Institute of Technology graduate students under the direction of the CSDL staff. DSST was developed with an emphasis on accuracy and computational efficiency. The semi-analytical theory has been used extensively to study the long-term evolution of orbits. The constants in the semi-analytical theory, the mean orbital elements, have also proven quite useful as the solve for variables in both batch least squares and recursive filter orbit determination processes based on the semi-analytical theory. The historical evolution of DSST can be seen in Reference².

Nonlinear Dynamics Web Tools, Nondy^{Web Tools 10}, is an e-Science project being developed in the University of La Rioja. The aim of this project is to build a non-commercial infrastructure to encourage scientific collaboration in which specialized software tools may be used freely through Internet. This project should promote not only e-Collaboration among the Web-Site's users, but also optimize use of resources, both human

and material. This project is not a closed one, but open to the collaboration of whoever wishes to participate in this initiative. Astrody^{Web Tools 11} is the specialized part related to Astrodynamics and Celestial Mechanics.

A brief description of the DSST standalone characteristic is given in this paper. Then the Astrody^{Web Tools} framework is presented. Finally we describe the integration of DSST into the Web-site and show its use through the Internet.

II. DSST

DSST^{12,13,14,15,16,17,18} is based on a semi-analytical satellite theory expressed in nonsingular equinoctial elements^{19,20}, which allows taking advantage of the accuracy of Special Perturbations (numerical integration) and the efficiency of General Perturbations (analytical satellite theory) because the influence of the long and short-period perturbations can be decoupled using the Perturbation Theory, and thus handled separately. Note that the equinoctial elements are related to the classical Keplerian elements by the following equations:

$$\begin{aligned}
 a &= a \\
 h &= e \sin(\omega + \Omega) \\
 k &= e \cos(\omega + \Omega) \\
 p &= \tan(i/2) \sin(\Omega) \\
 q &= \tan(i/2) \cos(\Omega) \\
 \lambda &= M + \omega + \Omega
 \end{aligned}
 \tag{1}$$

The equinoctial elements given in Eq. (1) represent one possible choice for the nonsingular state variables²¹.

The Generalized Method of Averaging leads to the Lagrangian VOP form given in Eq.(2) which is useful for perturbations with a conservative potential and the Gaussian VOP form (Eq.(4) in Ref³).

$$\frac{d\bar{a}_i}{dt} = \sum_{j=1}^6 (\bar{a}_i, \bar{a}_j) \frac{\partial}{\partial \bar{a}_j} \left[\frac{1}{2\pi} \int_0^{2\pi} R dM \right]
 \tag{2}$$

The Gaussian VOP is very useful for atmospheric drag and solar radiation pressure perturbations where it may be difficult to express the RHS's of the equations of motion in orbital elements.

Large stepsizes are possible for both the Lagrangian and Gaussian VOP forms. The Gaussian VOP may be uniquely able to take advantage of parallel processing concepts.

The atmosphere drag and solar radiation pressure models referenced in Tables 1 and 2 employ the Gaussian VOP.

The statement “DSST computes the short periodic motion using Finite Fourier series” applies to both Lagrangian VOP and Gaussian VOP.

Force models currently considered in DSST for the mean element equations of motion and for short periodic motion are shown in Tables 1 and 2. Refinement of the J_2 -squared model to better address high eccentricity cases is currently being investigated by Z. Folcik and P. Cefola.

Perturbation factors	Mean element equations of motion
Zonal harmonics including C_{20}	Linear terms in general form
Second-degree zonal harmonic	Second-order terms for the a, h, k, p, q, λ rates Terms of order $J_2^2 e^2$ neglected
Tesseral 1m harmonics of geopotential ($2 < l < 50, 1 < m < 50$)	Linear terms, including resonance effects in general form. Modified expansion for the Hansen coefficients
Attraction of the Moon and the Sun	Linear terms in general form
Atmosphere drag	Linear and cross with C_{20} terms. Rates evaluated via quadratures (Harris-Priester, Jacchia-Roberts ²² , MSISE-90)
Solar pressure	Linear terms of direct solar pressure. Rates evaluated via quadratures. Cylindrical model for shadow
Solid Earth Tides	Love number term

Table 1: DSST mean element equation of motion

DSST also includes a semi-analytical theory for the partial derivatives of perturbed motion and an efficient interpolation strategy, which greatly assists in producing the perturbed position and velocity and the partial derivatives at output request times for a given satellite.

GTDS R&D is an orbit determination system with orbit propagators, observation sensor models, and nonlinear estimation technology. DSST is one of the several orbit propagators that are available within GTDS. Other orbit propagators available within GTDS include Cowell numerical integration (fixed time step), Cowell numerical integration (time regularized), Brouwer, Brouwer-Lyddane, NORAD GP Theories, and NAVSPASUR PPT2.

The Cowell orbit propagators are general with respect to the physical models. Brouwer, Brouwer-Lyddane, the NORAD GPs, and NAVSPASUR PPT2 all have similar physical models: secular and long-periodic motion due to J_2 through J_4 and short-periodic motion due to J_2 . The DSST includes physical modeling that is nearly as general as Cowell.

Perturbation factors	Short periodic motion
Zonal harmonics including C_{20}	First-order terms in a, h, k, p, q, λ treated via a closed-form expansion in true longitude
Second-degree zonal harmonic	Second-order terms in a, h, k, p, q, λ treated via an expansion in true longitude Terms of order $J_2^2 e^2$ neglected
Tesseral 1m harmonics of geopotential ($2 < l < 50, 1 < m < 50$)	Linear terms in general form – partitioned into three categories 1. ‘m-dailies’ 2. linear-combination terms 3. J_2 /tesseral m-daily coupling terms
Attraction of the Moon and the Sun	Linear terms in general form. Closed form expansion in eccentric longitude. Weak time dependent terms
Atmosphere drag	Linear terms via an expansion in mean longitude. Coefficients evaluates via quadratures
Solar pressure	Linear terms via an expansion in mean longitude. Coefficients evaluated via quadratures
Solid Earth Tides	No

Table 2: DSST short periodic motion

III. ASTRODYNAMICS WEB TOOLS

The current prototype of Astrody^{Web}_{Tools} allows that through a user-friendly web interface, the user chooses the application, introduces the data and selects the appropriate constraints in an intuitive and easy way, with the help of the options in the graphical interface. After that, the application is executed in real-time and the critical information about the program behavior and

the output (graphical representation of the data, statistical analysis or any manipulation thereof whatsoever) are shown via the same web interface or downloaded to his computer.

Astrodynamics Web Tools

HOME SOFTWARE REPORTS USER AREA LINK STAFF CONTACT

Astrodynamics Tools

Universidad de La Rioja Dr. Juan Félix San Juan (juanfelix.sanjuan@unirioja.es)

Orbit Propagator Programs

Twelve mathematical zonal and tesseral models for prediction of satellite position and velocity using state vector are available:

Model	Coefficient	Order	Name	Real-Time
Zonal	J_2	2	ppkbJ2or2	Yes
	J_2	3	ppkbJ2or3	Yes
	J_2	4	ppkbJ2or4	No
	$J_2 \dots J_4$	2	ppkbJ4or2	No
	$J_2 \dots J_4$	3	ppkbJ4or3	No
	$J_2 \dots J_6$	2	ppkbJ6or2 ₍₁₎	No
	$J_2 \dots J_6$	3	ppkbJ6or3 ₍₁₎	No
	$J_2 \dots J_8$	2	ppkbJ8or2	No
Tesseral	2 x 2	4	tes2x2	Yes
	4 x 4	4	tes4x4	No
	6 x 6	4	tes6x6	No
	8 x 8	4	tes8x8	No

Real Observatorio de la Armada Dr. Martín Lara (mlara@roa.es)

ZERGOF: Zonal Earth Repeat Ground-track Orbits Finder

ZERGOF₍₂₎ is an application intended to help artificial satellite mission designers in their search for repeated ground-track, frozen orbits.

Dr. Paul Cefola (paulcefola@buffalo.edu)

Draper Semianalytical Satellite Theory (DSST) Standalone

DSST Standalone is an efficient Orbit Propagator based on a Semi-analytical Satellite theory which allows the determination of the orbits of artificial satellites and space debris objects.

Real Observatorio de la Armada y Universidad de La Rioja Dr. Martín Lara (mlara@roa.es)
Dr. Juan Félix San Juan (juanfelix.sanjuan@unirioja.es)

Repeat Ground-track Orbits Finder

It is an application intended to help artificial satellite mission designers in their search for repeated ground-track, frozen orbits in the case of planetary satellite.

1. Previous versions of these models were installed at the Centre National D'Etudes Spatiales (CNES).
2. The ZERGOF code used is a translation to ANSI C by J. F. San Juan although the original code was implemented in FORTRAN by M. Lara.

Fig. 1: Available applications in Astrody^{Web}_{Tools}

Figure 1 shows the four current applications available in Astrody^{Web}_{Tools}. Besides DSST, the others application are:

- Orbit Propagator Programs^{23,24} is an application, which contains twelve Analytical Orbit Propagator Programs (AOPP). These orbit propagators calculate the orbiter's position and velocity at any given moment directly by means of a function of time and the initial position and velocity of the orbiter. Every AOPP has been automatically obtained from an analytical approximation to the artificial satellite problem using a code generator.

The AOPPs have been classified first by gravity coefficients (i.e. zonal or tesseral), then by the selected coefficients and finally by the accuracy of the analytical approximation used.

- ZERGOF²⁵ (Zonal Earth Repeat Ground-track Orbits Finder) is a software package designed to search for repeating ground-track orbits automatically, in the case of the Earth, in a proportion between the number of nodal periods of the satellite on its orbit and the number of nodal days previously selected by the user. These kinds of orbits are highly desirable as nominal orbits for a variety of missions for artificial satellites, because their eccentricity and argument of the perigee remain almost constant or *frozen* during a long period of time.
- Repeat Ground-track Orbits Finder is a software package designed to search for repeating ground-track orbits automatically in the case of a planetary satellite. The search for an orbit of this kind begins when an approximate solution is obtained by means of analytical techniques. Subsequently, the above approximate solution is refined by using numerical continuation methods. The first part of this process is made using a *Mathematica* package, whereas the second is available in our Web-Site at this moment.




Fig. 2: Basic architecture of Astrody^{Web}_{Tools}

Figure 2 shows an overview of the architecture proposed for Astrody^{Web}_{Tools}. Through our framework the registered user can select and execute one of the available applications after completing the appropriate form with the initial values and parameters. The user-data are converted into the input of the application, which is executed, and the results are stored in files. After that, the user-results are processed by other open source applications such as Gnuplot, Octave, the statistical package R, Latex, or any other needed applications, and their outputs are embedded in the web

page. In addition, the graphics, pdf reports, and other outputs, can be downloaded directly by the user in several formats —for instance, the images can be obtained in eps or pdf format.

For example, in the case of ZERGOFF, the search for a frozen orbit begins when the user selects its characteristics. The gravity model, the number of zonal harmonics, the repeat ground track constrain and the type of periodic orbit, which is classified in near-circular Sun-Synchronicity orbit, near-circular with a selected inclination orbit, critical-inclination Sun-Synchronicity with argument of the perigee 90^0 or 270^0 orbit, or critical-inclination with an argument of the perigee 90^0 or 270^0 and a selected eccentricity orbit. The application obtains an approximate solution using analytical techniques. Subsequently, this solution is refined by means of numerical continuation methods.




Fig. 3: ZERGOFF Web interface

Figure 3 shows an example of the use of ZERGOFF. This searches the initial conditions of an almost circular orbit with a 56^0 orbit inclination, where 5 revolutions in 3 days is the repeat ground track constraint. The model only considers between J_2 and J_9 zonal harmonics.




Fig. 4: ZERGOFF output

Figure 4 shows the results provided by the Web-site after the execution of the application. The links in the left part of the interface facilitate graphical displays of the obtained results.

The trace of the orbit can be seen in Figure 5, whilst Figure 6 shows the evolution of the long-period terms.




Fig. 5: Satellite trace




Fig. 6: Long-period terms ($e \sin g, e \cos g$)

IV. USER DSST THROUGH ASTRODYNAMICS WEB TOOLS

DSST is a Fortran 77 based application, which does not have any type of graphical user interface (GUI), thus its execution is made from a command line. Currently, the DSST standalone source code is maintained by Z. Folcik and P Cefola. Figure 7 shows the basic DSST execution. The application reads the physical model files, which contain:

- astrodynamics constants data
- Solar/Lunar/Planetary ephemerides in either the J2000 or true data coordinates
- time conversion coefficients and polar motion coefficients
- 50x50 geopotential models
- quasi-logarithmic planetary geomagnetic indices and night-time minimum exospheric temperatures for the Jacchia-Roberts atmospheric density model

and the user-data files, which contain the initial conditions of the satellite and the configuration parameters of DSST and the output times selected by the user. Once data and constants are stored in memory, DSST carries out the calculus and the outputs are stored in two files.




Fig. 7: DSST executed mode

This executed mode represents a classical pattern in scientific programming, which we could call a black-box paradigm. The application is treated like a black box, as users only need to modify some input data files so as to run the application without re-compiling each time. Our Web-site provides these applications with an easy-to-use graphical-user web-interface which simplifies communication with the applications, such that their integration into the Astrody^{Web}_{Tools} framework is almost immediate, and then the details of their implementation, programming languages used, and so on, can be relegated to a secondary plane.

The current DSST Web interface allows the user to introduce the epoch, epoch mean elements set (Keplerian or Equinoctial), dynamical parameters, and so on, in two basic ways: the user-data file can be uploaded to Astrody^{Web}_{Tools}, as Figure 8 shows, or the user can fill in a web form, as Figure 9 shows.

Fig. 8: Upload user-data file

Figure 10 shows the results provided by the Web-site after the execution of DSST. The information shown is obtained from the two output files generated by this application: dsst.output and SPGOUT. The first contains the output data. The second is a text file, which stores some of the DSST force model parameters used in its execution and some of its intermediate output: element rate and short period model partial derivative

parameters, as well as short-periodic model options and short-periodic Fourier coefficients.

Fig. 9: DSST Web form




Fig. 10: DSST Web form

V. CONCLUSION AND FUTURE WORKS

The creation of an open source software suite of basic SSA tools available to all space actors is an important step towards enhancing safe and efficient operations in space. While the project is being developed, one of the tools considered, DSST standalone, can be used through Internet. DSST is an accurate, efficient and extensively used Semi-Analytical Satellite Theory. Astrody^{Web}_{Tools} is the Web framework, which provides DSST with a friendly GUI and users with a web service through which they can easily access using a Web browser.

In our project, DSST users play a relevant role in the improvement of this Web service, which is generated from the DSST output, because it is possible to modify the DSST Web interface in function of users' needs. For example, download files in a specific format, generate all kinds of graphics, allow uploading a file with their data in order to compare with the DSST output, or whatever users need. This collaboration intends to maximize the full utility of DSST output.

We are developing a new web environment, which will replace the current prototype. This will be available by the end of 2011 and Web 2.0 facilities and e-Learning tools will be added, in order to organize courses centered on these applications for those researchers and students who are interested in learning the theoretical knowledge they are based on.

VI. ACKNOWLEDGMENTS

Part of this research has been supported by the Government of Spain (Projects AYA 2009-11896, AYA 2010-18796, and grant Gobierno de La Rioja Fomenta 2010/16). The work of Zachary Folcik was sponsored by the U. S. Air Force under Air Force Contract FA8721-05-C-0002. Opinions, interpretations, conclusions, and recommendations are those of the authors and are not necessarily endorsed by the United States Government. Paul Cefola's effort was partially supported by the Secure World Foundation.

¹ L. K. Newman, R. Frigm, and D. McKinley, It's Not a Big Sky After All: Justification for a Close Approach Prediction and Risk Assessment Process, AAS 2009-369, presented at the AAS/AIAA Astrodynamics Specialist Conference, Pittsburgh, PA, 9-13 August 2009 (Vol. 135, Advances in the Astronautical Sciences)

² P. J. Cefola, B. Weeden and C. Levit, Open Source Software Suite for Space Situational Awareness and Space Object Catalog Work, 4th International Conference on Astrodynamics Tools Techniques, Madrid, Spain, 3-6 May 2010

³ P. J. Cefola, D. Phillion, and K. S. Kim, Improving Access to the Semi-Analytical Satellite Theory, AAS 09-341, presented at the AAS/AIAA Astrodynamics Specialist Conference, Pittsburgh, PA, August 2009

⁴ D. Brouwer, Solution of the Problem of Artificial Satellite Theory Without Drag, Astronomical Journal, Vol. 64, No. 1274, 1959, pp. 378-397

⁵ F. R. Hoots and R. L. Roehrich, Models for Propagation of NORAD Element Sets, Spacetrack Report No. 3, December 1980, Aerospace Defense Command, USAF

⁶ F. R. Hoots and R. G. France, An Analytic Satellite Theory Using Gravity and a Dynamic Atmosphere, Celestial Mechanics, Vol. 40, No. 1, 1987, pp. 1-18

⁷ PPT2: The NAVSPASUR Model of Satellite Motion, NAVSPASUR Report 92-01, July 1992

⁸ P. J. Cefola and D. J. Fonte, Extension of the Naval Space Command Satellite Theory PPT2 to include a General Tesseral M-daily Model, pre-print AIAA 96-3606, presented at the AIAA/AAS Astrodynamics Specialist Conference, San Diego, CA, August 1996

⁹ V. F. Boikov, G. N. Makhonin, A. V. Testov, Z. N. Khutorovsky, and A. N. Shogin, Prediction Procedures Used in Satellite Catalog Maintenance, Journal of Guidance, Control, and Dynamics, Vol. 32, No. 4, 2009, pp. 1179-1199

¹⁰ J. F. San Juan, R. López and I. Pérez, Nondy^{Web}_{Tools} : Nonlinear Dynamics Web Tools, 4th International Conference on Astrodynamics Tools Techniques, Madrid, Spain, 3-6 May 2010

¹¹ J. F. San Juan, R. López. L. M. López and I. Pérez, Symbolic-numeric cooperation in an open science project, Symbolic and Numeric Algorithms for Scientific Computing (IEEE Computer Society), 2011, pp. 37-46

¹² W. D. McClain, A Recursively Formulated First-Order Semianalytic Artificial Satellite Theory Based on the Generalized Method of Averaging, Vol. 1 (The Generalized Method of Averaging Applied to the Artificial Satellite Problem), Computer Sciences Corporation CSC/TR-77/6010, 1977 (an updated version of this document is available from W. D. McClain at the Draper Laboratory).[the 'blue book']

¹³ W. D. McClain, A Recursively Formulated First-Order Semianalytic Artificial Satellite Theory Based on the Generalized Method of Averaging, Vol. 2 (The Explicit Development of the First-Order Averaged Equations of Motion for the Nonspherical Gravitational and Nonresonant Third-Body Perturbations), Computer Sciences Corporation CSC/TR-78/6001, 1978 [the 'yellow book']

¹⁴ L. W. Early, A Portable Orbit Generator Using Semianalytical Satellite Theory, AIAA Paper 86- 2164-CP, AIAA/AAS Astrodynamics Conference, Williamsburg, VA, August 1986

¹⁵ R. J. Proulx, Mathematical Description of the Tesserel Resonance and Resonant Harmonic Coefficient Solve-For Capabilities, Draper Laboratory Intralab Memorandum NSWC-001-15Z-RJP, 13 April 1982

¹⁶ J. G. Neelon, P. J. Cefola, and R. J. Proulx, Current Development of the Draper Semianalytical Satellite Theory Standalone Orbit Propagator Package", AAS Pre-print 97-731, presented at the AAS/AIAA Astrodynamics Conference, Sun Valley, ID, August 1997

¹⁷ D. A. Danielson, B. Neta, and L. W. Early, Semianalytic Satellite Theory (SST): Mathematical Algorithms, Naval Postgraduate School, NPS Report NPS-MA-94-001, January 1994

¹⁸ M. Slutsky, Zonal Harmonic Short-Periodic Model Developed for the Precision Orbit Propagation (POP) Contract, Draper Laboratory Intralab Memorandum PL-016-81-MS, 30 November 1981

¹⁹ P. J. Cefola, Equinoctial Orbit Elements – Application to Artificial Satellite Orbits, AIAA 72-937, presented at the 1972 AIAA/AAS Astrodynamics Specialist Conference, Palo Alto, CA, September 1972

²⁰ P. J. Cefola, A Recursive Formulation for the Tesserel Disturbing Function in Equinoctial Variables, AIAA paper 76-839, presented at the AIAA/AAS Astrodynamics Specialist Conference, San Diego, CA, August 1976

²¹ G. R. Hintz, Survey of Orbit Element Sets, Journal of Guidance, Control, and Dynamics, Vol. 31, No. 3, 2008, pp. 785-790

²² Charles E. Roberts, An Analytic Model for Upper Atmosphere Densities Based Upon Jacchia's 1970 Models, Celestial Mechanics Vol. 4(3,4), 1971, pp. 368-378

²³ J. F. San Juan, ATESAT: Automatization of theories and ephemeris in the artificial satellite problem, CT/TI/MS/MN/94-250, 1994, CNES, France

²⁴ J. F. San Juan and S. Serrano, Application of the Z6PPKB ATESAT-model to compute the orbit of an artificial satellite around Mars, DTS/MPI/MS/MN/2000-057, 2000, CNES, France

²⁵ M. Lara, Searching for Repeating Ground Track Orbits: A Systematic Approach, The Journal of the Astronautical Sciences, Vol. 47, 1999, 177-188